For Research Use Only

CoraLite® Plus 488 Anti-Human CD81 (5A6) Mouse IgG2a Recombinant Antibody



Catalog Number: CL488-65644

Basic Information

Catalog Number: CL488-65644

Size:

100tests, 5 ul/test

Source: Mouse

Isotype: IgG2a

GenBank Accession Number:

BC002978 GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000110651

Full Name: CD81 molecule Calculated MW:

26 kDa

Purification Method:

Protein A purification

CloneNo.: 5A6

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 491 nm / 516 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human peripheral blood leukocytes,

Background Information


CD81 (also known as TAPA1 or TSPAN28) is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Many of these members are expressed on leukocytes and have been implicated in signal transduction, cell-cell interactions, and cellular activation and development. CD81 is involved in signal transduction and cell adhesion in the immune system (PMID: 9597125). CD81 has also been identified as en essential receptor for HCV (hepatitis C virus) (PMID: 21428934).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 human peripheral blood leukocytes were surface stained with 5 ul Coralite® Plus 488 Anti-Human CD81 (5A6) Mouse IgG2a RecAb (CL488-65644, Clone: 5A6) or Coralite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.



1x10^6 human peripheral blood leukocytes were surface stained with 5 ul Coralite® Plus 488 Anti-Human CD81 (5A6) Mouse IgG2a RecAb (CL488-65644, Clone: 5A6) (red) or Coralite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were not fixed.