For Research Use Only

CoraLite® Plus 488-conjugated RBAP48 Monoclonal antibody

Catalog Number: CL488-66060

Featured Product



Basic Information

Catalog Number: GenBank Accession Number: CL488-66060 BC053904

Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 ug/ml by 5928 Nanodrop: **UNIPROT ID:** Recommended Dilutions:

Q09028 Mouse Full Name:

Isotype retinoblastoma binding protein 4

IgG2b Calculated MW:

Immunogen Catalog Number: 48 kDa AG6196

Observed MW:

53 kDa

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human

5C4D6

IF/ICC 1:50-1:500

Purification Method:

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Background Information

Histone-binding protein RBBP4 (also known as RbAp48, or NURF55) is a protein that in humans is encoded by the RBBP4 gene. This gene encodes a ubiquitously expressed nuclear protein that belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2/NuRD complex complex that has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of corepressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. A decrease of RbAp48 in the dentate gyrus (DG) of the hippocampus in the brain is suspected to be a main cause of memory loss in normal aging (PMID: 23986399).

Positive Controls:

IF/ICC: A549 cells,

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A549 cells using CoraLite® Plus 488-conjugated RBAP48 antibody (CL488-66060, Clone: 5C4D6) at dilution of 1:200, CL594-Phalloidin (red).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human RBAP48 (CL488-66060, Clone:5C4D6) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).