CoraLite® Plus 488-conjugated Fetuin-A/AHSG Monoclonal antibody

Catalog Number: CL488-66094


Basic Information	Catalog Number: CL488-66094	GenBank Accession Number: BC048198	Purification Method: Protein A purification				
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG2a Immunogen Catalog Number: AG10073	GenelD (NCBI): 197	CloneNo.: 1F6B9				
		UNIPROT ID: P02765 Full Name: alpha-2-HS-glycoprotein Calculated MW: 367 aa, 39 kDa	Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima				
				Observed MW: 58 kDa			
				Applications	Tested Applications:	Positive Controls:	
					IF/ICC, FC (Intra) Species Specificity: human	IF/ICC : H	epG2 cells,
		Background Information		Fetuin-A (alpha-2-HS-glycoprotein, AHSG), a liver borne plasma protein, contributes to the prevention of soft tissue calcification, modulates inflammation, reduces INS sensitivity and fosters weight gain following high fat diet or ageing. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. The protein is commonly present in the cortical plate of the immature cerebral cortex and bone marrow hemopoietic matrix, and it has therefore been postulated that it participates in the development of the tissues.			
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage						

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Antibodies | ELISA kits | Proteins www.ptglab.com

Selected Validation Data





Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 Fetuin-A antibody (CL488-66094, Clone: 1F6B9) at dilution of 1:200.

1x10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human Fetuin-A (CL488-66094, Clone:1F6B9)(red), or 0.8 ug CoraLite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).