For Research Use Only

CoraLite® Plus 488-conjugated MPO Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-66177

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-66177 BC130476 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 4353

Nanodrop: **UNIPROT ID:** P05164

Mouse Full Name: Isotype: myeloperoxidase IgA Calculated MW:

Immunogen Catalog Number: 745 aa, 84 kDa AG17564 Observed MW:

90 kDa

Purification Method:

Thiophilic affinity chromatograph

CloneNo.: 4C11F6

Recommended Dilutions:

IF-P 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications: IF-P, FC (Intra) Species Specificity:

human, rat

Positive Controls:

IF-P: human tonsillitis tissue,

Background Information

The MPO gene encodes myeloperoxidase, a lysosomal hemoprotein located in the azurophilic granules of polymorphonuclear (PMN) leukocytes and monocytes. In response to stimulation, MPO is activated into a transient intermediate with potent antimicrobial oxidizing abilities (PMID: 17650507). The mRNA is translated into a single protein of 90 kDa, which displays enzymatic activity and undergoes proteolytic maturation into a heavy chain of 59 kDa and a light chain of 13.5 kDa; these subunits then dimerize into the mature tetramer and the mature MPO is a heterotetramer composed of two identical heavy chains and two identical light chains (PMID:12773517). The 24-kDa material had a map identical to that of 13.5 kDa subunit and represents a dimer of the 13.5 kDa subunit (PMID:3008892). Defects in MPO are the cause of myeloperoxidase deficiency (MPOD). It has 3 isoforms produced by alternative splicing.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using Coralite® Plus 488 MPO antibody (CL488-66177, Clone: 4C11F6) at dilution of 1:200.



1X10^6 HL-60 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human MPO (CL488-66177, Clone:4C11F6) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).