For Research Use Only

## CoraLite® Plus 488-conjugated P62/SQSTM1 Monoclonal antibody



Catalog Number: CL488-66184

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-66184 BC017222 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 8878

Nanodrop: **UNIPROT ID:** Q13501

Mouse Full Name: Isotype: sequestosome 1

IgG2a Calculated MW:

Immunogen Catalog Number: 48 kDa

AG13131 Observed MW:

62 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 1H5C1

Recommended Dilutions:

IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity:

human

Positive Controls:

IF/ICC: HepG2 cells, Chloroquine treated U2OS cells

## **Background Information**

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and sometimes aggregated proteins for autophagic degradation. As a autophagy-specific substrate, p62 is degraded during the autophagic process, which makes intracellular level of p62 as a marker for autophagy flux. p62 is at the cross-roads of several signaling pathways including Ras/ Raf/ MAPK and NFkB and plays important role in cancer. p62 is a component of inclusion bodies/ protein aggregates found in human diseases, including Huntington's disease, Alzheimer's disease, Parkinson's disease in the brain, and nephropathic cystinosis in kidney (PMID: 22074114, 22860231, 22714671). The molecular weight of p62 is predicted as 48/38 kDa, while western blot analyses using this antibody demonstrate the major band around 60-62 kDa in various tissues.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HepG2 cells using CL488-66184 (P62,SQSTM1 antibody) at dilution of 1:100.



 $Immun of luorescent analysis of (-20 ^{\circ}C Ethanol) fixed HepG2 cells using CL488-66184 (P62,SQSTM1 antibody) at dilution of 1:100.$