For Research Use Only

CoraLite® Plus 488-conjugated PD-1/CD279 Monoclonal antibody



Catalog Number: CL488-66220

Basic Information

Catalog Number: GenBank Accession Number:

CL488-66220 BC074740 GeneID (NCBI):

100ul, Concentration: 2000 µg/ml by 5133

Source

programmed cell death 1 Mouse Calculated MW:

Isotype: 288 aa, 32 kDa lgG2b

Immunogen Catalog Number:

AG12470

Purification Method: Protein A purification

CloneNo.: 4H4D1

Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 488 nm / 515 nm

Applications

Tested Applications:

FC. IF

Species Specificity: human, rat, mouse

Positive Controls:

IF: human tonsillitis tissue,

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436). The calculated molecular weight of PD-1 is 32 kDa. It has been reported that PD-1 is heavily glycosylated and migrates with an apparent molecular mass of 47-55 kDa on SDS-PAGE (PMID: 8671665; 17640856; 17003438). This anitbody is CL488(Ex/Em 488 nm/515 nm) conjugated.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using the CoraLite® Plus 488-conjugated version of this antibody, CL488-66220 (PD-1/CD279 antibody), at dilution of 1:150.



1X10^6 Jurkat cells untreated (blue, dashed line) or treated with CD3 antibody (OKT3, 2 ug/ml) were surface stained with 0.4 ug CoraLite® Plus 488 Anti-Human PD-1/CD279 (CL488-66220, Clone: 4H4D1) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue, solid line). Cells were not fixed.