For Research Use Only

## CoraLite® Plus 488-conjugated IL-12/IL-23 p40 Monoclonal antibody



**Purification Method:** 

Protein A purification

Excitation/Emission maxima

CloneNo.:

wavelengths: 493 nm / 522 nm

4B2A8

Catalog Number: CL488-66288

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-66288 BC067498 GeneID (NCBI):

100ul , Concentration: 1000  $\mu g/ml$  by 3593 Nanodrop: **UNIPROT ID:** 

P29460 Mouse Full Name:

Isotype: interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic IgG2a lymphocyte maturation factor 2, p40)

Immunogen Catalog Number: Calculated MW: AG21429

328 aa, 37 kDa

Observed MW: 37 kDa

**Applications** 

**Tested Applications:** 

Species Specificity: human, pig

## **Background Information**

IL-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed  $of the \ IL-12\ p40\ subunit\ linked\ to\ the\ IL-12\ p35\ subunit,\ and\ the\ heterodimer\ signals\ through\ the\ IL-12\ receptor\ (IL-12\ p40\ subunit\ linked\ to\ the\ large p40\ subunit\ linked\ to\ the\ large p40\ subunit\ linked\ to\ the\ large p40\ subunit\ linked\ to\ the\ large\ p40\ subunit\ large\ p40\ subuni$ 12R), which comprises the IL-12R $\beta$ 1 and IL-12R $\beta$ 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R $\beta$ 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data