For Research Use Only

CoraLite® Plus 488-conjugated Cyclin D3 Monoclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL488-66357

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-66357 BC011616

Size: GeneID (NCBI): 100ul , Concentration: 1000 ug/ml by 896

 Nanodrop;
 ENSEMBL Gene ID:

 Source:
 ENSG00000112576

 Mouse
 UNIPROT ID:

 Isotype:
 P30281

 IgG1
 Full Name:

Immunogen Catalog Number:cyclin D3AG22825Calculated MW:33 kDa

Observed MW: 33-35 kDa Purification Method:

Protein G purification

CloneNo.: 1F2C6

Recommended Dilutions: IF/ICC 1:300-1:1200

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human

Positive Controls:

IF/ICC: HeLa cells,

Background Information

Cyclin D3 belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation.

Storage

Storage:


Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite® Plus 488 Cyclin D3 antibody (CL488-66357, Clone: 1F2C6) at dilution of 1:600.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Cyclin D3 (CL488-66357, Clone:1F2C6) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).