For Research Use Only

CoraLite® Plus 488-conjugated Beta Catenin Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-66379

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** CL488-66379 NM 001904 Protein G purification

> GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 ug/ml by 1499 1B8A1

Nanodrop: ENSEMBL Gene ID: Recommended Dilutions: Source: ENSG00000168036 IF/ICC 1:50-1:500

Mouse UNIPROT ID: Excitation/Emission maxima

Isotype: P35222 wavelengths: 493 nm / 522 nm lgG1 Full Name:

catenin (cadherin-associated protein),

beta 1, 88kDa Calculated MW: 781 aa, 86 kDa

Applications

Tested Applications: Positive Controls: IF/ICC, FC (Intra) IF/ICC: MCF-7 cells,

Species Specificity: human, mouse, rat, pig

Background Information

β-Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β-Catenin was originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to a-catenin and the actin cytoskeleton. Besides its essential role in the AJs, β -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ β -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β -catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222) This antibody is conjugated with CL488(Ex/Em 488 nm/515 nm).

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CoraLite® Plus 488 Beta Catenin antibody (CL488-66379, Clone: 188A1) at dilution of 1:200, Alpha Tubulin antibody (11224-1-AP, red).



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Beta Catenin (CL488-66379, Clone:1B8A1) (red), or 0.4 ug CoraLite® Plus 488 Mouse [gG1 lsotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).