For Research Use Only

CoraLite® Plus 488-conjugated SIRT2 Monoclonal antibody



Purification Method:

Catalog Number: CL488-66410

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-66410 BC003547 Protein G purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 ug/ml by 22933 1D8G10 Nanodrop:

UNIPROT ID: Recommended Dilutions: Q8IXJ6 IF/ICC 1:50-1:500

Mouse Full Name: Excitation/Emission maxima

Isotype: sirtuin (silent mating type wavelengths: information regulation 2 homolog) 2 493 nm / 522 nm lgG1

(S. cerevisiae) Immunogen Catalog Number:

Calculated MW: AG7756 43 kDa Observed MW: 37-45 kDa

Applications

Tested Applications:

IF/ICC

Species Specificity: human, mouse, rat, pig **Positive Controls:**

IF/ICC: HepG2 cells,

Background Information

The silent information regulator(SIR2) family of genes are highly conserved from prokaryotes to eukaryotes and are involved in diverse processes, including transcriptional regulation, cell cycle progression, DNA-damage repair and aging. SIR2 contains a 323 amino acid catalytic core domain with a NAD-binding domain and a large groove which is the likely site of catalysis. SIR2 is widely expressed, highly expressed in heart, brain and skeletal muscle, while it is weakly expressed in placenta and lung. Down-regulated in many gliomas suggesting that it may act as a tumor $suppressor\ gene\ in\ human\ gliomas\ possibly\ through\ the\ regulation\ of\ microtubule\ network.\ SIRT2\ exists\ various$ isoforms and the molecular weight of isoforms are 35kDa, 40 kDa, and 42 kDa.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

E: proteintech@ptglab.com

W: ptglab.com

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL488-66410 (SIRT2 antibody) at dilution of 1:100.