For Research Use Only

# CoraLite® Plus 488-conjugated TXN Monoclonal antibody



**Purification Method:** 

Catalog Number: CL488-66475

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-66475 BC003377 Protein A purification
Size: GeneI D (NCBI): CloneNo.:

Size: Genel D (NCBI): CloneNo 100ul , Concentration: 1000 µg/ml by 7295 1F3C1

Nanodrop; Full Name: Recommended Dilutions:
Source: thioredoxin IF 1:500-1:2000

Mouse Calculated MW: Excitation/Emission maxima Isotype: 12 kDa wavelengths:

IgG2a Observed MW: 488 nm / 515 nm

IgG2a Observed MW:

Immunogen Catalog Number: 12 kDa

AG6355

Applications

**Tested Applications:** 

IF

**Species Specificity:** 

Human

Positive Controls:

IF: HepG2 cells,

### **Background Information**

TXN, TRDX, TRX, TRX1, ADF and SASP, belongs to the thioredoxin family. It participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. TXN plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. TXN induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity.

#### Storage

#### Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488-conjugated TXN antibody (CL488-66475, Clone: 1F3C1) at dilution of 1:1000, C594-Phalloidin (red).