## For Research Use Only

## CoraLite® Plus 488-conjugated TXN Monoclonal antibody



Catalog Number: CL488-66475

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-66475 GeneID (NCBI): Size:

100ul, Concentration: 1000 ug/ml by 7295 Nanodrop; **UNIPROT ID:** 

Source P10599 Mouse Full Name: Isotype thioredoxin

IgG2a Calculated MW: Immunogen Catalog Number: 12 kDa

AG6355 Observed MW:

12 kDa

**Tested Applications:** 

IF/ICC

**Purification Method:** 

Protein A purification

CloneNo.: 1F3C1

**Recommended Dilutions:** IF/ICC 1:500-1:2000

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

Species Specificity:

Human

**Positive Controls:** 

IF/ICC: HepG2 cells,

## **Background Information**

TXN, TRDX, TRX1, ADF and SASP, belongs to the thioredoxin family. It participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. TXN plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. TXN induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488-conjugated TXN antibody (CL488-66475, Clone: 1F3C1) at dilution of 1:1000, C594-Phalloidin (red).