For Research Use Only

CoraLite® Plus 488-conjugated OGT Monoclonal antibody

Catalog Number:CL488-66823 Featured Product




Basic Information	Catalog Number: CL488-66823	GenBank Accession Number: BC014434	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG2a	GenelD (NCBI): 8473	CloneNo.: 2B2A6
		UNIPROT ID: 015294	Recommended Dilutions: WB 1:1000-1:4000
		Full Name: O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N- acetylglucosamine:polypeptide-N- acetylglucosaminyl transferase)	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	Calculated MW: 1046 aa, 117 kDa		
	Observed MW: 110 kDa		
	Applications	Tested Applications: WB	Positive Controls:
WB : HepG2 cells, Species Specificity: human, mouse, rat, pig			
Background Information	O-linked N-acetylglucosamine transferase (OGT) catalyzes the attachment of N-acetylglucosamine (GlcNAc) monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression. O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL488-66823 (OGT antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.