For Research Use Only

CoraLite® Plus 488-conjugated NeuN Monoclonal antibody

Catalog Number:CL488-66836 2 Publications




Basic Information	Catalog Number: CL488-66836	GenBank Accession Number: NM_001082575		Purification Method: Protein G purification
	Size: 100ul , Concentration: 1000 ug/ml by	Full Name: Recommended Dilutions hexaribonucleotide binding protein 3 IF-P 1:50-1:500		
	Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number:			Excitation/Emission maxima wavelengths:
Applications	AG28016 Tested Applications:	Positive Controls:		
	IF-P, FC (Intra) Cited Applications: IF Species Specificity: human, mouse, rat Cited Species: mouse, rat		IF-P : mouse o	erebellum tissue, mouse brain tissue
Background Information	NeuN, encoded by FOX3, is a neuron-specific nuclear protein. Anti-NeuN stains exclusively neuronal cells in the central and peripheral nervous systems, especially postmitotic and differentiating neurons, as well as terminally differentiated neurons. Anti-NeuN has been used widely as a reliable tool to detect most postmitotic neuronal cell types. The immunohistochemical staining is primarily localized in the nucleus of the neurons with lighter staining in the cytoplasm.			
Notable Publications	Author Pu	bmed ID	Journal	Application
	Jingcheng Shi 38	585980	bioRxiv	
	Lirong Zhang 36	801237	Exp Eye Res	IF
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Procli Aliquoting is unnecessary for -20°C s	n300, 0.5% BSA,		


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using CoraLite® Plus 488 NeuN antibody (CL488-66836, Clone: 3A4C1) at dilution of 1:200.



1X10^6 U-87 MG cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human NeuN (CL488-66836, Clone:3A4C1) (red), or 0.4 ug Mouse IgG1 Isotype Control (66360-1-Ig, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



1X10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human NeuN (CL488-66836, Clone:3A4C1) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).