For Research Use Only

## CoraLite® Plus 488-conjugated XRN2 Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-66852

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-66852 BC006417 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 22803 Nanodrop: **UNIPROT ID:** Q9H0D6

Mouse Full Name: Isotype: 5'-3' exoribonuclease 2

lgG1 Calculated MW: Immunogen Catalog Number: 104 kDa AG27927 Observed MW:

109 kDa

**Purification Method:** Protein G purification

CloneNo.:

2C3E3

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: Human, mouse, rat

## **Background Information**

XRN2 is one exonuclease that degrades the Pol II associated product of poly(A) site cleavage, which is crucial for Pol II termination. During transcription termination, XRN2 cleavages at the polyadenylation site liberates a 5' fragment which is subsequently processed to form the mature mRNA and a 3' fragment which remains attached to the elongating polymerase. The processive degradation of this 3' fragment by this protein may promote termination of transcription.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human XRN2 (CL488-66852, Clone:2C3E3) (red), or 0.8 ug CoraLite® Plus 488 Mouse IgG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).