For Research Use Only

CoraLite® Plus 488-conjugated ATX3,ATXN3 Monoclonal antibody

www.ptglab.com

Purification Method:

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

wavelengths: 493 nm / 522 nm

CloneNo.:

1F7E10

Catalog Number: CL488-67057

Basic Information

Catalog Number: GenBank Accession Number:

CL488-67057 BC033711 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 4287

Nanodrop: **UNIPROT ID:** P54252

Mouse Full Name: Isotype: ataxin 3

IgG2a Calculated MW: Immunogen Catalog Number: 370 aa, 43 kDa

AG6627 Observed MW: 41-45 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse, pig Positive Controls:

IF/ICC: A549 cells,

Background Information

ATXN3, which has deubiquitinase activity and act as a component of the ubiquitin proteasome system, plays a role in transcriptional regulation and neuroprotection. ATXN3 interacts with RAD23, HHR23A and HHR23B, involves in the pathology of MJD. ATXN3 is a mixed-linkage, chain-editing enzyme and that the UIM region of ATXN3 regulates its substrate specificity. Contains an N-terminal deubiquitinating domain, called the Josephin domain, followed by 2 ubiquitin-interacting motifs (UIMs) and a polyQ tract near the C terminus. ATXN3 can be phosphorylated in a protein casein kinase-2-dependent manner, thus the MW would be larger than the predicted one.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A549 cells using Coralite® Plus 488 ATX3,ATXN3 antibody (CL488-67057, Clone: 1F7E10) at dilution of 1:200, CL594-phalloidin (red).



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human ATX3,ATXN3 (CL488-67057, Clone:1F7E10) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL488-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).