For Research Use Only

CoraLite® Plus 488-conjugated MCM4 Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-67103

Basic Information

Catalog Number: GenBank Accession Number:

CL488-67103 BC031061 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 4173 Nanodrop: **UNIPROT ID:**

P33991 Mouse Full Name:

Isotype: minichromosome maintenance lgG2b complex component 4

Immunogen Catalog Number: Calculated MW: AG28642 863 aa. 97 kDa

> Observed MW: 97 kDa

Applications

Tested Applications:

Species Specificity: human, mouse, rat

Purification Method: Protein A purification

CloneNo.: 2H2A1

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Background Information

DNA replication licensing factor MCM4 (MCM4), also named Cdc21, acts as component of the MCM2-7 complex (MCM $complex) which is the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replication in the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, are the putative replica$ elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity.

Positive Controls: IF/ICC: U2OS cells,

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using CoraLite® Plus 488 MCM4 antibody (CL488-67103, Clone: 2H2A1) at dilution of 1:200.