For Research Use Only

## CoraLite® Plus 488-conjugated ACC1 Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-67373

Featured Product

1 Publications

**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

CL488-67373

BC137287 GeneID (NCBI):

CloneNo.:

100ul, Concentration: 1000 ug/ml by 31

1A11G10

Nanodrop:

**UNIPROT ID:** Q13085

Recommended Dilutions:

493 nm / 522 nm

Mouse

Full Name:

IF/ICC 1:50-1:500

Isotype:

acetyl-Coenzyme A carboxylase

Excitation/Emission maxima wavelengths:

IgG2a

alpha

Immunogen Catalog Number: AG17503

Calculated MW:

2383 aa, 275 kDa

Observed MW:

260-270 kDa

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Positive Controls: IF/ICC: HeLa cells,

Species Specificity: human, mouse, rat

**Background Information** 

ACACA(Acetyl-CoA carboxylase 1, ACC), also named as ACAC, ACC1 and ACCA, belongs to the biotin containing enzyme family. It catalyzes the synthesis of malonyl-CoA, which is an intermediate substrate playing a pivotal role in the regulation of fatty acid metabolism and energy production. ACACA is involved in the biosynthesis of fatty acids, and malonyl-CoA produced is used as a building block to extend the chain length of fatty acids by fatty acid synthase (FAS)(PMID:19900410). It has 4 isoforms produced by alternative promoter usage with the molecular weight between 260 kDa and 270 kDa.

**Notable Publications** 

Author **Pubmed ID** Application Journal Victor Gray 39488214 Cell Metab

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 ACC1 antibody (CL488-67373, Clone: 1A11G10 ) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human ACC1 (CL488-67373, Clone:1A11G10) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL488-6360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).