For Research Use Only

## CoraLite® Plus 488-conjugated RIG-1/DDX58 Monoclonal antibody



**Purification Method:** 

Recommended Dilutions:

Excitation/Emission maxima

IF 1:50-1:500

wavelengths:

488 nm / 515 nm

Catalog Number: CL488-67556

**Basic Information** 

Catalog Number: GenBank Accession Number:

 CL488-67556
 BC132786
 Protein G purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ul , Concentration: 1000 μg/ml by 23586
 3B10F5

Nanodrop; Full Name:

Source: DEAD (Asp-Glu-Ala-Asp) box

Mouse polypeptide 58

Isotype: Calculated MW:
IgG1 925 aa, 107 kDa

Immunogen Catalog Number: Observed MW: AG18591 101/106 kDa

**Applications** 

**Tested Applications:** 

IF

Species Specificity: Human, mouse, rat **Positive Controls:** 

IF: HepG2 cells,

## **Background Information**

DDX58, also named as RIG-1, belongs to the helicase family. It is involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAV5/IP51, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). Detects dsRNA produced from non-self dsDNA by RNA polymerase III, such as Epstein-Barr virus-encoded RNAs (EBERs). It is essential for the production of interferons in response to RNA viruses including paramyxoviruses, influenza viruses, Japanese encephalitis virus and HCV. The antibody is specific to DDX58,which has two isoform with the MW of 106 and 101 kDa.

Storage

Storage:

Store at -20°C. Avoid exposure to light.


Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Coralite® Plus 488-conjugated RIG-1/DDX58 antibody (CL488-67556, Clone: 3B10F5) at dilution of 1:100.