For Research Use Only

CoraLite® Plus 488-conjugated EIF4A3 Monoclonal antibody

www.ptglab.com

Purification Method:

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

493 nm / 522 nm

wavelengths:

CloneNo.:

1G6H9

Catalog Number: CL488-67740

Basic Information

Catalog Number: GenBank Accession Number:

CL488-67740 BC003662 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 9775 Nanodrop: **UNIPROT ID:** P38919

Mouse Full Name:

Isotype: eukaryotic translation initiation

lgG2b factor 4A, isoform 3 Immunogen Catalog Number: Calculated MW:

AG11130 47 kDa

Observed MW: 47 kDa

Applications

Tested Applications:

Species Specificity: Human, mouse

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

EIF4A3 is a component of the exon junction complex (EJC), which assembles near exon-exon junctions of mRNAs as a result of splicing. EJC proteins involves in postsplicing events, including mRNA export, cytoplasmic localization, and nonsense-mediated decay. Its RNA-dependent ATPase and RNA-helicase activities are induced by CASC3, but abolished in presence of the MAGOH/RBM8A heterodimer, thereby trapping the ATP-bound EJC core onto spliced mRNA in a stable conformation. Besides, it involved in translational enhancement of spliced mRNAs after formation of the 805 ribosome complex and binds spliced mRNA in sequence-independent manner, 20-24 nucleotides upstream of mRNA exon-exon junctions


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 EIF4A3 antibody (CL488-67740, Clone: 1G6H9) at dilution of 1:200.