For Research Use Only

CoraLite® Plus 488-conjugated Phospho-mTOR (Ser2448) Monoclonal antibody



Catalog Number: CL488-67778

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: Protein A purification

Size: GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 ug/ml by 2475 2A12G3

Nanodrop; Full Name: Recommended Dilutions:

Source: FK506 binding protein 12-rapamycin IF/ICC 1:50-1:500

 Mouse
 associated protein 1
 Excitation/Emission maxima

 Isotype:
 Calculated MW:
 wavelengths:

IgG2b 289 kDa 493 nm / 522 nm

Applications

Tested Applications:

IF/ICC, FC (Intra)

Positive Controls:

IF/ICC : HepG2 cells,

Species Specificity: human, mouse

Background Information

MTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTOR is phosphorylated at Ser2448 via the PI3 kinase/Akt signaling pathway and autophosphorylated at Ser2481. mTOR plays a key role in cell growth and homeostasis and may be abnormally regulated in tumors.

Storage Stora


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 PhosphomTOR (Ser2448) antibody (CL488-67778, Clone: 2A12G3) at dilution of 1:100. DAPI (blue)



1X10^6 Calyculin A treated HeLa cells were intracellularly stained with 0.25 ug CoraLite® Plus 488 Anti-Human Phospho-mTOR (Ser2448) (CL488-67778, Clone:2A12G3) (red), or 0.25 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 90% MeOH.