

For Research Use Only

CoraLite® Plus 488-conjugated EIF4A2 Monoclonal antibody

Catalog Number: **CL488-67799**



Basic Information

Catalog Number: CL488-67799	GenBank Accession Number: BC012547	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 1974	CloneNo.: 2D11E11
Source: Mouse	UNIPROT ID: Q14240	Recommended Dilutions: IF/ICC 1:50-1:500
Isotype: IgG2a	Full Name: eukaryotic translation initiation factor 4A, isoform 2	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Immunogen Catalog Number: AG9085	Calculated MW: 408 aa, 46 kDa	
	Observed MW: 46 kDa	

Applications

Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : HEK-293 cells,
Species Specificity: Human, Rat, Mouse	

Background Information

Eukaryotic initiation factor 4A (EIF4A) has an essential role in the binding of mRNA to the 43S preinitiation complex when protein synthesis begins, as EIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon. EIF4A2 is a ATP-dependent RNA helicase, which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome.


Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using Coralite® Plus 488 EIF4A2 antibody (CL488-67799, Clone: 2D11E11) at dilution of 1:100.



1X10⁶ HEK-293 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human EIF4A2 (CL488-67799, Clone:2D11E11) (red), or 0.4 ug Coralite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).