For Research Use Only

CoraLite® Plus 488-conjugated MYDGF/C19orf10 Monoclonal antibody



493 nm / 522 nm

Catalog Number: CL488-68131

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** CL488-68131 BC010129 Protein G purification

> GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 µg/ml by 56005 1H8F4

Full Name: Recommended Dilutions:

Source: chromosome 19 open reading frame IF 1:50-1:500

Mouse

Excitation/Emission maxima Isotype: Calculated MW: wavelengths:

173 aa, 19 kDa lgG1

Immunogen Catalog Number: Observed MW: 15-19 kDa AG13438

Applications Tested Applications: Positive Controls:

Human, Mouse, Rat

FC (Intra), IF IF: HepG2 cells, Species Specificity:

Background Information

Myeloid-derived growth factor (MYDGF), also named as chromosome 19 open reading frame 10 (C19orf10), is an $FLS-derived\ protein\ that\ is\ secreted\ into\ the\ synovial\ fluid.\ MYDGF\ as\ a\ paracrine-acting\ protein\ secreted\ from$ monocytes/macrophages that improves tissue repair and heart function after myocardial infarction. MYDGF has been reported to regulates neutrophil motility in interstitial tissue damage(PMID: 34047769).

Storage Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Coralite® Plus 488 MYDGF/C19orf10 antibody (CL488-68131, Clone: 1H8F4) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human MYDGF/C19orf10 (CL488-68131, Clone:1H8F4) (red), or 0.8 ug CoraLite® Plus 488 Mouse I gG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).