For Research Use Only

## CoraLite® Plus 488-conjugated TPMT Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-68251

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-68251 BC005339 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7172

Nanodrop: **UNIPROT ID:** P51580 Mouse

Full Name: Isotype: thiopurine S-methyltransferase

IgG2a Calculated MW:

Immunogen Catalog Number:

AG28693 Observed MW:

28 kDa

32 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

human

**Purification Method:** Protein A purification

CloneNo.: 1D4B3

Recommended Dilutions:

IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

IF/ICC

Positive Controls:

IF/ICC: HepG2 cells,

## **Background Information**

The TPMT gene is localized to chromosome 6p22.3 and is approximately 34 kDa in length. It has an open reading frame (ORF) of 735 bp that encodes for a 245-amino acid peptide. The wild type TPMT allele, expressing the enzyme with normal activity. To date, there have been 30 variant TPMT alleles described (PMID: 22493988). Thiopurine compounds are commonly used in the treatment of childhood acute lymphoblastic leukemia, and as immunosuppressants following organ transplantation or for treatment of various autoimmune disorders. Thiopurine S-methyltransferase (TPMT) is required for detoxification, through S-methylation, of 6-thioguanine nucleotides (TGNs), a byproduct of thiopurine metabolism. Catalog#10682-1-AP is a rabbit polyclonal antibody raised against the full-length of human TPMT. TPMT has a predicted molecular weight of 28 kDa, however, as a result of its phosphorylation it separates on a SDS gel as a band of approximately 28-32 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 TPMT antibody (CL488-68251, Clone: 1D4B3) at dilution of 1:200, CL594-Phalloidin (red).