

For Research Use Only

CoraLite®488-conjugated Phospho-Beta Catenin (Ser33) Recombinant antibody



Catalog Number:CL488-80067

Basic Information

Catalog Number: CL488-80067	GenBank Accession Number: BC058926	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 1499	CloneNo.: 3K1
Source: Rabbit	Full Name: catenin (cadherin-associated protein), beta 1, 88kDa	Excitation/Emission maxima wavelengths: 488 nm/515 nm
Isotype: IgG	Calculated MW: 781 aa, 86 kDa	
	Observed MW: 90 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
Human

Background Information

β-Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β-Catenin was originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to α-catenin and the actin cytoskeleton. Besides its essential role in the AJs, β-catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/β-catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β-catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222). CK1 phosphorylates β-Catenin at Ser45. This phosphorylation event primes β-Catenin for subsequent phosphorylation by GSK-3β. GSK-3β destabilizes β-catenin by phosphorylating it at Ser33, Ser37, and Thr41. Mutations at these sites result in the stabilization of β-Catenin protein levels and have been found in many tumor cell lines .

Storage

Storage:
Store at -20°C. Avoid exposure to light.


Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ PC-3 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug CoraLite® 488 Anti-Human Phospho-Beta Catenin (Ser33) (CL488-80067, Clone:3K1), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.