For Research Use Only

## CoraLite® Plus 488-conjugated Ecadherin Recombinant antibody

Catalog Number: CL488-80541-7



**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-80541-7 BC141838 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 999

Nanodrop: **UNIPROT ID:** P12830 Rabbit Full Name:

Isotype: cadherin 1, type 1, E-cadherin (epithelial)

IgG Immunogen Catalog Number: Calculated MW: 882 aa, 97 kDa AG14973

Observed MW: 135 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

**Purification Method:** 

Protein A purification

CloneNo.: 240159A1

Recommended Dilutions:

IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

## **Background Information**

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. E-cadherin (epithelial cadherin), also known as CDH1 (cadherin 1) or CAM 120/80, is a classical member of the cadherin superfamily which also include N-, P-, R-, and B-cadherins. E-cadherin is expressed on the cell surface in most epithelial tissues. The extracellular region of E-cadherin establishes calcium-dependent homophilic trans binding, providing specific interaction with adjacent cells, while the cytoplasmic domain is connected to the actin cytoskeleton through the interaction with  $p120-, a-, \beta-, and \gamma-catenin (plakoglobin). E-cadherin is important in the maintenance of the epithelial integrity, and properties of the epithelial integrity and properties of the epithelial int$ is involved in mechanisms regulating proliferation, differentiation, and survival of epithelial cell. E-cadherin may also play a role in tumorigenesis. It is considered to be an invasion suppressor protein and its loss is an indicator of high tumor aggressiveness.

Positive Controls: IF/ICC: MCF-7 cells,


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Coralite® Plus 488 E-cadherin antibody (CL488-80541-7, Clone: 240159A1) at dilution of 1:200.