## For Research Use Only

# CoraLite® Plus 488-conjugated AKT1-Specific Recombinant antibody

Catalog Number:CL488-80816 Featured Product

## **Basic Information**

Catalog Number: CL488-80816	GenBank Accession Number: BC 000479
Size: 100ul , Concentration: 1000 ug/ml by	GeneID (NCBI): 207
Nanodrop;	UNIPROT ID:
Source:	P31749
Rabbit	Full Name:
lsotype: IgG	v-akt murine thymoma viral oncogene homolog 1
Immunogen Catalog Number:	Calculated MW:
AG0213	56 kDa
	Observed MW:

#### **Purification Method:** Protein A purification CloneNo.: 509 Excitation/Emission maxima wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** FC (Intra) Species Specificity: human, mouse, rat

### **Background Information**

The serine-threonine protein kinase AKT1 is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcriptionindependent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery.

#### **Storage**

#### Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage


56-62 kDa

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



# Selected Validation Data



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human AKT1 (CL488-80816, Clone:509) (red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000, Clone: ) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).