For Research Use Only

CoraLite® Plus 488-conjugated ENO1/2/3 Recombinant antibody



Purification Method:

wavelengths:

Positive Controls:

493 nm / 522 nm

Catalog Number: CL488-81504

Basic Information

Catalog Number: GenBank Accession Number:

CL488-81504 BC015641 Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 µg/ml by 2023 6F24

Recommended Dilutions: Source: IF 1:200-1:800 enolase 1, (alpha) Rabbit Calculated MW: Excitation/Emission maxima

Isotype: 47 kDa IgG Observed MW:

Immunogen Catalog Number: 47 kDa

Applications

Tested Applications:

IF: HepG2 cells,

Species Specificity: Human, Mouse, Rat, Pig

Background Information

Enolase (ENO), also known as 2-phospho-D-glycerate hydrolase, is a metalloenzyme that catalyzes the dehydration of 2-phospho-D-glycerate into phosphoenolpyruvate in the glycolytic pathway. There are three different ENO isoforms in higher eukaryotes, including ENO1 or a-ENO, ENO2 or γ -ENO, and ENO3 or β -ENO. Furthermore, the expression of the different ENO isoforms is tissue specificity. ENO1 is expressed in almost all kinds of tissue, whereas ENO3 is restricted to muscle tissues and ENO2 is described as a neuron- and neuroendocrine tissue-specific enolase. ENO 1 serves multifunctional roles in physiological and pathological processes. (PMID: 32256815). This antibody can recognize ENO1, ENO2, and ENO3.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 ENO 1/2/3 antibody (CL488-81504, Clone: 6E24) at dilution of 1:400.