For Research Use Only

## CoraLite® Plus 488-conjugated IBA1 Recombinant antibody

www.ptglab.com

**Purification Method:** 

CloneNo.:

IF-P 1:50-1:500

wavelengths:

493 nm / 522 nm

Protein A purification

Excitation/Emission maxima

Catalog Number: CL488-81728

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-81728 GeneID (NCBI): Size:

100ul , Concentration: 1000 ug/ml by 199 4D5 Nanodrop; **ENSEMBL Gene ID:** Recommended Dilutions:

Source ENSG00000204472 Rabbit **UNIPROT ID:** 

Isotype P55008 IgG **Full Name:** 

Immunogen Catalog Number: allograft inflammatory factor 1

AG1363 Calculated MW:

17 kDa Observed MW: 17 kDa

**Applications** 

**Tested Applications:** IF-P, FC (Intra)

Species Specificity: Human, mouse, rat

**Positive Controls:** 

IF-P: rat brain tissue,

## **Background Information**

IBA1 is a 143 amino acid cytoplasmic, inflammation response scaffold protein. It is constitutively expressed in monocytes and macrophages and is known to be involved in macrophage activation. It is a marker of activated macrophage. Expression of IBA1 is up-regulated in activated microglia following facial nerve axotomy, ischemia, and several brain diseases, thereby implicating it in the activated phenotypes of microglia.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)


## **Selected Validation Data**



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Coralite® Plus 488 IBA1 antibody (CL488-81728, Clone: 4D5) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using CoraLite® Plus 488 IBA1 antibody (CL488-81728, Clone: 4D5) at dilution of 1:200



1X10^6 THP-1 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human IBA1 (CL488-81728, Clone:4D5) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).