For Research Use Only

CoraLite® Plus 488-conjugated PRDX3 Recombinant antibody

Catalog Number:CL488-81833 Featured Product

Basic Information	Catalog Number: CL488-81833	GenBank Accession Number: BC007062	Purification Method: Protein A purification	
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1062	GeneID (NCBI): 10935	CloneNo.: 1L11	
		UNIPROT ID: P30048	Recommended Dilutions: IF/ICC 1:50-1:500	
		Full Name: peroxiredoxin 3 Calculated MW: 27 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
				Observed MW: 28 kDa
		Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls:
IF/ICC : HepG2 cells, Species Specificity: human, mouse, rat			epG2 cells,	
Background Information	PRDX3(Peroxiredoxin 3), also named as AOP1, HBC189 and MER5, belongs to the ahpC/TSA family. It is involved in redox regulation of the cell and protects radical-sensitive enzymes from oxidative damage by a radical-generating system. PRDX3 is required for MYC-mediated proliferation, transformation, and apoptosis after glucose withdrawal and essential for maintaining mitochondrial mass and membrane potential in transformed rat and human cells (PMID:12011429). PRDX3 acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol.			
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin		nt.	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


proteintech

Antibodies | ELISA kits | Proteins www.ptglab.com

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 PRDX3 antibody (CL488-81833, Clone: 1L11) at dilution of 1:200.



1x10^6 MCF-7 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human PRDX3 (CL488-81833, Clone:1L11)(red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).