For Research Use Only

CoraLite® Plus 488-conjugated STK39 Recombinant antibody

www.ptglab.com

Catalog Number: CL488-83081-2

Basic Information

Catalog Number: GenBank Accession Number:

CL488-83081-2 BC166614 GeneID (NCBI): CloneNo.: Size:

230382G9 100ul , Concentration: 1000 ug/ml by 27347 Nanodrop; **UNIPROT ID:**

Q9UEW8 Source Rabbit Full Name:

Isotype serine threonine kinase 39

IgG (STE20/SPS1 homolog, yeast)

Immunogen Catalog Number: Calculated MW:

AG34503 59 kDa Observed MW:

62 kDa

Purification Method:

Protein A purification

Recommended Dilutions:

IF-P 1:50-1:500 IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

IF/ICC, IF-P IF-P: mouse brain tissue. Species Specificity: IF/ICC: NIH/3T3 cells. human, rat

Background Information

STK39 (Serine/Threonine Kinase 39) is a proline- and alanine-rich Ste20-related kinase (also named as SPAK), which is composed of a short N-terminal proline and alanine repeats (PAPA box), a kinase catalytic domain and a Cterminal regulatory domain (PMID: 16382158). In mammalians, STK39 plays an important role in ion homeostasis by regulating the cation chloride cotransporters' activities, which is crucial for the modulation of renal salt transport $and blood\ pressure\ (PMID:\ 33500714).\ Notably,\ STK39\ is\ also\ implicated\ in\ the\ regulation\ of\ tumor\ cell\ proliferation,$ migration, and invasion in multiple cancers, including osteosarcoma and cervical cancer (PMID: 34335956).

Positive Controls:

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using CoraLite® Plus 488 STK39 antibody (CL488-83081-2, Clone: 230382G9) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Coralite® Plus 488 STK39 antibody (CL488-83081-2, Clone: 230382G9) at dilution of 1:200.



1x10^6 Daudi cells were intracellularly stained with 0.8 ug Coralite® Plus 488 Stk39 Recombinant Antibody (CL488-83081-2, Clone:230382G9)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).