For Research Use Only

CoraLite® Plus 488-conjugated OGT Recombinant antibody

Catalog Number: CL488-83113



Purification Method:

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

CloneNo.:

230385D11

IF/ICC 1:50-1:500

493 nm / 522 nm

wavelengths:

Basic Information

Catalog Number: GenBank Accession Number:

CL488-83113 BC014434 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 8473 Nanodrop: **UNIPROT ID:** Source 015294

Rabbit Full Name:

Isotype: O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N-IgG

acetylglucosamine:polypeptide-N-Immunogen Catalog Number: acetylglucosaminyl transferase) AG2160

Calculated MW: 1046 aa, 117 kDa

Observed MW: 110 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: Hela cells,

Background Information

O-linked N-acetylglucosamine transferase (OGT) catalyzes the attachment of N-acetylglucosamine (GlcNAc) monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression, O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed Hela cells using CoraLite® Plus 488 OGT antibody (CL488-83113, Clone: 230385D11) at dilution of 1:200.



1x10^6 HeLa cells were intracellularly stained with 0.8 ug Coralite® Plus 488 OGT Recombinant Antibody (CL488-83113, Clone:230385D11)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).