For Research Use Only

CoraLite® Plus 488-conjugated BAG3 Recombinant monoclonal antibody

www.ptglab.com

Catalog Number: CL488-83779-4

Basic Information

Catalog Number: GenBank Accession Number:

CL488-83779-4 BC006418 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 9531 Nanodrop: **UNIPROT ID:** 095817

Rabbit Full Name:

Isotype: BCL2-associated athanogene 3 IgG Calculated MW:

Immunogen Catalog Number: 61 kDa

AG0956 Observed MW:

74-80 kDa

Purification Method:

Protein A purification

CloneNo.: 240860F12

Recommended Dilutions:

IF/ICC: 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

IF/ICC

Species Specificity:

human

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

BAG3 (Bcl2-associated athanogene 3) belongs to the BAG protein family, the co-chaperone that binds to Hsc70/Hsp70 through the BAG domain and modulates their activity in polypeptide folding. BAG3 contains also a WW domain and a proline-rich (PXXP) repeat, that mediate binding to partners different from Hsp70. Through interacting with different molecular partner, BAG3 influences several cell processes, such as apoptosis, autophagy and cell motility. BAG3 protein has been reported to sustain cell survival, resistance to therapy, and/or motility and metastatization in several tumor types, thus being identified as a potential target for anticancer therapies. In addition, defects in BAG3 are the cause of some myopathy. BAG3 normally migrates around 74-80 kDa; a slightly different molecular weight or a doublet form can be observed in some cell types and/or following cell exposure to stressors. A synaptosome associated form of 40 kDa has recently been described.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Coralite® Plus 488 BAG3 antibody (CL488-83779-4, Clone: 240860F12) at dilution of 1:200.