For Research Use Only

CoraLite® Plus 488 Anti-Human IFNgamma Rabbit Recombinant Antibody

Catalog Number: CL488-98187



Basic Information

Catalog Number:

BC070256

Purification Method: Protein A purification

CL488-98187 Size:

GeneID (NCBI):

CloneNo.:

100tests, 5 ul/test

3458

241390B10

Source: Rabbit

ENSEMBL Gene ID: ENSG00000111537

GenBank Accession Number:

Excitation/Emission maxima wavelengths:

Isotype:

UNIPROT ID: P01579

493 nm / 522 nm

Immunogen Catalog Number: EG0004

Full Name: IFN gamma

Calculated MW:

166 aa, 19 kDa

Applications

Tested Applications:

FC (Intra)

human

Positive Controls:

Species Specificity:

FC (Intra): PMA, Ionomycin and Brefeldin A treated

human PBMCs.

Background Information

Interferon gamma (IFN γ), is a type II interferon that provides immunity against bacterial, viral and protozoan infections. In its active form, IFN γ is a glycosylated, non-covalently linked homodimer of 29-32 kDa subunits. It is produced by a number of immune cell types including natural killer cells, natural killer T cells, and effector lymphocyte T cells following antigenic and inflammatory triggers. The IFN γ dimer binds to its cognate receptor which has two subunits: IFN-y R1 which is the ligand-binding chain (a chain) and IFN-y R2, the signal-transducing chain (β chain). Binding to the receptor activates the JAK/STAT pathway which in turn activates IFN γ responsive genes. While IFN γ can inhibit viral replication, it also works as an immune-modulator and immune-stimulator by increasing surface expression of class I MHC proteins. (PMID: 19268625; 10688427)


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 untreated or PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with 5 ul Coralite® Plus 488 Anti-Human IFN-gamma Rabbit RecAb (CL488-98187, Clone:241390810). Cells were co-stained with Coralite® Plus 647 Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (CL647-65569, Clone: OKT3). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with 5 ul Coralite® Plus 488 Anti-Human IFN-gamma Rabbit RecAb (CL488-98187, Clone: 241390B10) (red) or Coralite® Plus 488 Rabbit IgG Isotype Control RecAb (CL488-98136, Clone: 240953C9)(blue).

1x10^6 untreated human PBMCs were intracellularly stained with 5 ul Coralite® Plus 488 Anti-Human IFN-gamma Rabbit RecAb (CL488-98187, Clone: 241390B10)

