For Research Use Only

CoraLite®555-conjugated S100A11 Polyclonal antibody



Catalog Number: CL555-10237

Featured Product

Basic Information

Catalog Number:

CL555-10237 BC001410 GeneID (NCBI):

100ul, Concentration: 500 ug/ml by 6282

Nanodrop:

UNIPROT ID: P31949 Rabbit Full Name:

Isotype S100 calcium binding protein A11

IgG Calculated MW: Immunogen Catalog Number: 105 aa, 12 kDa

AG0343

Tested Applications: IF/ICC, FC (Intra)

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: PC-3 cells,

Background Information

S100A11 (also known as S100C or calziggarin), is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs (PMID: 18694925). S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. First discovered in 1989, S100A11 has since been proposed to have direct biological functions in an assortment of physiological processes such as endo- and exocytosis, regulation of enzyme activity, cell growth and regulation, apoptosis and inflammation (PMID: 15241500). Chromosomal rearrangements and altered expression of ${\tt S100A11\,have\,been\,implicated\,in\,tumor\,metastasis.\,This\,S100A11\,antibody\,(10237-1-AP)\,has\,also\,been\,implicated\,in\,tumor\,metastasis.}$ instrumental in investigations into S100A11's role in the development of brain tumors in tuberous sclerosis complex (TSC) patients (PMID: 20133820).

GenBank Accession Number:

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using CoraLite®555 S100A11 antibody (CL555-10237) at dilution of 1:200.



1X10^6 PC-3 cells were intracellularly stained with 0.4 ug Coralite®555 Anti-Human S100A11 (CL555-10237) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).