### For Research Use Only

# CoraLite®555-conjugated c-MYC Polyclonal antibody

Catalog Number:CL555-10828 Featured Product



**Purification Method:** 

IF/ICC 1:50-1:500

wavelengths:

557 nm / 570 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL555-10828 BC000141
Size: GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 4609
Nanodrop; UNIPROT ID:
Source: P01106
Rabbit Full Name:

 Isotype:
 v-myc myelocytomatosis viral

 IgG
 oncogene homolog (avian)

Immunogen Catalog Number: Calculated MW:
AG1263 49 kDa

Observed MW: 62-65 kDa, 50 kDa

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse Positive Controls:

IF/ICC: HEK-293 cells,

#### **Background Information**

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC. The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

#### Storage

**Applications** 

#### Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using CoraLite®555 c-MYC antibody (CL555-10828) at dilution of 1:200.



1X10^6 Raji cells were intracellularly stained with 0.2 ug CoraLite® 555 Anti-Human c-MYC (CL555-10828) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).