For Research Use Only

CoraLite®555-conjugated P16-INK4A Polyclonal antibody Catalog Number:CL555-10883 Featured Product




Basic Information	Catalog Number: CL555-10883	GenBank Accession Number: BC021998	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1328	GenelD (NCBI): 1029	Recommended Dilutions: IF/ICC 1:50-1:500
		UNIPROT ID: P42771 Full Name: cyclin-dependent kinase inhibitor Calculated MW: 16 kDa Observed MW: 16-18 kDa	Excitation/Emission maxima wavelengths: 557 nm / 570 nm
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human	Positive Controls: IF/ICC : HeLa cells,	
Background Information	CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively-spliced variants encoding distinct proteins proteins were reported. Two of them named p16-INK4 and p14 are sharing 50% identity. The third one -p14(ARF) is entirely unrelated. 10883-1-AP reacts with p16 specifically. P16 plays an essential role in regulating the cell cycle, and mutations in p16 increase the risk of developing various cancers, including melanoma.		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	1300, 0.5% BSA, pH 7.3.	nt.


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®555 P16-INK4A antibody (CL555-10883) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite®555 Anti-Human P16-INK4A (CL555-10883) (red), or 0.8 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).