For Research Use Only

## CoraLite®555-conjugated Cyclin E2 Polyclonal antibody

www.ptglab.com

Catalog Number: CL555-11935

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL555-11935 BC020729 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 9134 Nanodrop: **UNIPROT ID:** 096020 Rabbit Full Name: Isotype: cyclin E2

IgG Calculated MW: Immunogen Catalog Number: 374 aa, 44 kDa AG2532 Observed MW: 44 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths: 557 nm / 570 nm

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity:

human

Positive Controls:

IF/ICC: HeLa cells,

## **Background Information**

Cyclin E2 (CCNE2) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of cyclindependent kinases (CDKs). Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of cell cycle events. CCNE2 forms a complex with and functions as a regulatory subunit of CDK2 and has been shown to specifically interact with CIP/KIP family of CDK inhibitors. CCNE2 plays a role in cell cycle G1/S transition and its expression peaks at the G1-S phase. Whereas cyclin E1 is expressed in most proliferating normal and tumor cells, cyclin E2 levels are low or undetectable in nontransformed cells, and are elevated in tumor-derived cells.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®555 Cyclin E2 antibody (CL555-11935) at dilution of 1:200.