For Research Use Only

CoraLite®555-conjugated Cytokeratin 19 Polyclonal antibody



Catalog Number: CL555-14965

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL555-14965 BC002539
Size: GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 3880Nanodrop;UNIPROT ID:Source:P08727RabbitFull Name:Isotype:keratin 19

Immunogen Catalog Number: 44 kDa

AG6830

human, mouse, rat

IgG

Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Purification Method:

Applications

Tested Applications: Positive Controls: IF/ICC, FC (Intra) IF/ICC: HeLa cells, Species Specificity:

Calculated MW:

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT19, one of type I keratins, is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. Due to its high sensitivity, KRT19 is the most used marker for detection of tumor cells disseminated in lymph nodes, peripheral blood, and bone marrow of breast cancer patients. This antibody, generated against full length KRT19 protein, also recognizes KRT17, another type I keratin homologous to KRT19.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite®555 Cytokeratin 19 antibody (CL555-14965) at dilution of 1:200.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Coralite®555 Anti-Human Cytokeratin 19 (CL555-14965) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).