For Research Use Only

CoraLite®555-conjugated HMMR-Specific Polyclonal antibody

www.ptglab.com

Catalog Number:CL555-15820

Featured Product

Basic Information

Catalog Number:

CL555-15820

Nanodrop:

Source:

Rabbit

IgG

GenBank Accession Number:

NM_001142556

GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 3161

UNIPROT ID:

075330

Full Name: Isotype: hyaluronan-mediated motility

receptor (RHAMM)

Calculated MW:

84 kDa

Observed MW:

70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity:

human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

Hyaluronan-mediated motility receptor (HMMR), also termed CD168, was first described by Turley in murine cells . It is reported that HMMR has an extensive coiled-coil structure (CC) that contains multiple sites for interactive partners. Initially, HMMR was considered a novel hyaluronan-mediated motility receptor and a microtubuleassociated spindle assembly factor. Full-length human RHAMM is an 85 kDa coiled-coil protein that occurs both in intracellular and extracellular compartments. It has highly restricted and tightly regulated expression in most normal tissues, but is one of a number of oncogenic proteins that are exported to the cell surface in response to tissue stress by unconventional transport mechanisms. (PMID: 36750558, PMID: 30249497)


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human HMMR-Specific (CL555-15820) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® 555 HMMR-Specific antibody (CL555-15820) at dilution of 1:200.