For Research Use Only

CoraLite®555-conjugated PI3 Kinase p110 Beta Polyclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Catalog Number: CL555-20584

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL555-20584 NM_006219 Antigen affinity purification
Size: Genel D (NCBI): Recommended Dilutions:

100ul , Concentration: 1000 μg/ml by 5291 IF 1:50-1:500

Nanodrop; Full Name: Excitation/Emission maxima

Source: phosphoinositide-3-kinase, catalytic, wavelengths:
Rabbit beta polypeptide 555 nm / 570 nm

Isotype: Calculated MW:
IgG 123 kDa
Observed MW:
110-130 kDa

Applications

Tested Applications: FC (Intra), IF

Species Specificity:

human, rat

Positive Controls:

IF: HeLa cells,

Background Information

PIK3CB(phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit beta isoform) is also named as PIK3C1, PI3K-beta, p110beta. The gene encodes a 1070 amino acid protein which belongs to the PI3/PI4-kinase family. Phosphoinositide 3-kinases (PI3Ks) have been implicated as participants in signaling pathways regulating cell growth by virtue of their activation in response to various mitogenic stimuli. The class I PI3 kinases are heterodimers composed of 110 kDa catalytic subunits that associate with regulatory adaptor proteins. Four class I catalytic subunits have been identified, PIK3CA (p110a), PIK3CB (p110β), PIK3CD (p110δ) and PIK3CG (p110γ) (PMID:19177002). This antibody is specific to PIK3CB.

Storage

Storage:

Store at -20°C. Avoid exposure to light.


Storage Buffe

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®555 Pl3 Kinase p110 Beta antibody (CL555-20584) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human PI3 Kinase p110 Beta (CL555-20584) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).