For Research Use Only

CoraLite®555-conjugated BCL10 Polyclonal antibody Catalog Number:CL555-50180 Featured Product




CL555-50180	BC053617	Antigen affinity purification
Size: 100ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG0037	GeneID (NCBI): 8915 UNIPROT ID: 095999 Full Name: B-cell CLL/lymphoma 10 Calculated MW: 233 aa, 26 kDa Observed MW: 26-32 kDa	Excitation/Emission maxima wavelengths: 557 nm / 570 nm
Applications IF/ICC, FC (Intra)		
Species Specificity: human, mouse, rat		
associated lymphoid tissue (MALT) ly	ymphoma. BCL10 contains a caspas	
Storage Buffer:	2	nt.
	100ul, Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG0037 Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat The gene encoding BCL10 was identi associated lymphoid tissue (MALT) by shown to induce apoptosis and to act Storage: Stora at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Procli	100ul, Concentration: 500 ug/ml by 8915 Nanodrop; UNIPROT ID: Source: 095999 Rabbit Full Name: Isotype: B-cell CLL/lymphoma 10 IgG Calculated MW: Immunogen Catalog Number: 233 aa, 26 kDa AG0037 Observed MW: 26-32 kDa Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat The gene encoding BCL10 was identified by its recurrent translocation to associated lymphoid tissue (MALT) lymphoma. BCL10 contains a caspase shown to induce apoptosis and to activate NF-kappaB. Storage: Storage: Storage: Storage to light. Stable for one year after shipme


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A549 cells using Coralite®555 BCL10 antibody (CL555-50180) at dilution of 1:0.



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human BCL10 (CL555-50180) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).