## For Research Use Only

## CoraLite® Plus 555 Anti-Human PD-1/CD279 (EH12.2H7)

Antibodies | ELISA kits | Proteins WWW.ptglab.com


Catalog Number: CL555-65168


Basic Information	Catalog Number: CL555-65168	GenBank Accession Number: BC074740	Purification Method: Affinity purification
	Size: 100tests , 5 ul/test	GenelD (NCBI): 5133	CloneNo.: EH12.2H7
	Source: Mouse Isotype: IgG1, kappa	UNIPROT ID: Q15116 Full Name: programmed cell death 1	Excitation/Emission maxima wavelengths: 554 nm / 570 nm
		Applications	
Species Specificity: Non-human primates, Human			
Background Information	Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).		
Storage	Storage: Store at 2-8°C. Avoid exposure to Storage Buffer: PBS with 0.09% sodium azide a	to light. Stable for one year after shipme nd 0.5% BSA.	nt.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





1X10^6 day 3 PHA treated human PBMCs were surface co-stained with CL750 Anti-Human CD3 and 5 ul Coralite® Plus 555 Anti-Human PD-1/CD279 (CL555-65168, Clone: EH12.2H7) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.

1X10<sup>^6</sup> day 3 PHA treated human PBMCs were surface stained with 5 ul Coralite® Plus 555 Anti-Human PD-1/CD279 (CL555-65168, Clone: EH12.2H7) (red) or Mouse IgG1 Isotype Control (blue). Unactivated PBMCs were surface stained with 5 ul Coralite® Plus 555 Anti-Human PD-1/CD279 (black). Cells were not fixed. Lymphocytes were gated.