For Research Use Only

## CoraLite® Plus 555 Anti-Human CD279 (EH12.2H7)



Catalog Number: CL555-65168

**Basic Information** 

**Applications** 

Catalog Number: CL555-65168

GenBank Accession Number: BC074740

**Purification Method:** Affinity purification

Size:

GeneID (NCBI):

288 aa, 32 kDa

CloneNo.: EH12.2H7

wavelengths:

100tests, 5 ul/test

5133 Full Name:

Excitation/Emission maxima

Mouse

programmed cell death 1 Calculated MW:

554 nm / 570 nm

Isotype: IgG1, kappa

**Tested Applications:** 

FC

Species Specificity:

Non-human primates, Human

**Background Information** 


Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

## Selected Validation Data



1X10^6 day 3 PHA treated human PBMCs were surface co-stained with CL750 Anti-Human CD3 and 5 ul Coralite® Plus 555 Anti-Human CD279 (CL555-65168, Clone: EH12.2H7) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 day 3 PHA treated human PBMCs were surface stained with 5 ul Coralite® Plus 555 Anti-Human CD279 (CL555-65168, Clone: EH12.2H7) (red) or Mouse IgG1 Isotype Control (blue). Unactivated PBMCs were surface stained with 5 ul Coralite® Plus 555 Anti-Human CD279 (black). Cells were not fixed. Lymphocytes were gated.