For Research Use Only

CoraLite®555-conjugated Lamin B1 Monoclonal antibody



Catalog Number: CL555-66095

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL555-66095 BC012295 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 4001

Nanodrop: ENSEMBL Gene ID: ENSG00000113368 Mouse

UNIPROT ID: Isotype: P20700 lgG1 Full Name: Immunogen Catalog Number: lamin B1 AG20522 Calculated MW:

> 66 kDa Observed MW:

66-70 kDa

Tested Applications:

Species Specificity:

human, mouse, rat

Purification Method:

Protein G purification

CloneNo.: 3C10G12

Recommended Dilutions: IF/ICC 1:500-1:2000

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

IF/ICC, FC (Intra)

Positive Controls:

IF/ICC: U2OS cells, HepG2 cells

Background Information

Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin. The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Vertebrate lamins consist of two types, A and B. This gene encodes one of the two B type proteins, B1. This protein is not suitable for samples where the nuclear envelope has been removed.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using CoraLite®555 Lamin B1 antibody (CL555-66095, Clone: 3C10G12) at dilution of 1:1000.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human Lamin B1 (CL555-66095, Clone:3C10G12) (red), or 0.4 ug isotype control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite@555 Lamin B1 antibody (CL555-66095, Clone: 3C 10G12) at dilution of 1:200.