For Research Use Only

CoraLite®555-conjugated Cytokeratin 18 Monoclonal antibody

www.ptglab.com

Catalog Number: CL555-66187

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** CL555-66187 BC000180 Protein A purification

> GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 ug/ml by 3875 1G11C4

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P05783 IF/ICC 1:50-1:500

Mouse Full Name: Excitation/Emission maxima wavelengths:

Isotype: keratin 18 557 nm / 570 nm lgG1 Calculated MW:

Immunogen Catalog Number: 48 kDa

AG1260

Applications Tested Applications: Positive Controls: IF/ICC, FC (Intra) IF/ICC: HepG2 cells,

Species Specificity:

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT18, also named as CYK18, PIG46 and K18, is the most commonly found members of the intermediate filament gene family. KRT18 is a type I keratin which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®555 Cytokeratin 18 antibody (CL555-66187, Clone: 1G11C4) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®555 Anti-Human Cytokeratin 18 (CL555-66187, Clone:1G11C4) (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).