

For Research Use Only

# CoraLite®555-conjugated PI3 Kinase p110 Beta Monoclonal antibody



Catalog Number:CL555-67121

## Basic Information

<b>Catalog Number:</b> CL555-67121	<b>GenBank Accession Number:</b> BC114432	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ul , Concentration: 1000 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5291	<b>CloneNo.:</b> 1B2A1
<b>Source:</b> Mouse	<b>Full Name:</b> phosphoinositide-3-kinase, catalytic, beta polypeptide	<b>Excitation/Emission maxima wavelengths:</b> 557 nm / 570 nm
<b>Isotype:</b> IgG1	<b>Calculated MW:</b> 1070 aa, 123 kDa	
<b>Immunogen Catalog Number:</b> AG17505	<b>Observed MW:</b> 120-130 kDa	

## Applications

**Tested Applications:**  
FC (Intra)  
**Species Specificity:**  
Human, Rat

## Background Information

PIK3CB(phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit beta isoform) is also named as PIK3C1, PI3K-beta, p110beta. The gene encodes a 1070 amino acid protein which belongs to the PI3/PI4-kinase family. Phosphoinositide 3-kinases (PI3Ks) have been implicated as participants in signaling pathways regulating cell growth by virtue of their activation in response to various mitogenic stimuli. The class I PI3 kinases are heterodimers composed of 110 kDa catalytic subunits that associate with regulatory adaptor proteins. Four class I catalytic subunits have been identified, PIK3CA (p110α), PIK3CB (p110β), PIK3CD (p110δ) and PIK3CG (p110γ) (PMID:19177002).

## Storage


**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
Aliquoting is unnecessary for -20°C storage

**\*\*\* 20ul sizes contain 0.1%BSA**

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human PI3 Kinase p110 Beta (CL555-67121, Clone:1B2A1) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).