For Research Use Only

CoraLite®555-conjugated STAT4 Monoclonal antibody

Catalog Number: CL555-67568



Purification Method:

IF/ICC 1:50-1:500

wavelengths: 557 nm / 570 nm

Excitation/Emission maxima

Protein G purification

Basic Information

Catalog Number: GenBank Accession Number:

CL555-67568 BC031212 GeneID (NCBI):

CloneNo.: 100ul, Concentration: 1000 ug/ml by 6775 4A8C9 Nanodrop: Recommended Dilutions:

UNIPROT ID: Q14765 Mouse

Full Name:

Isotype: signal transducer and activator of lgG1 transcription 4

Immunogen Catalog Number: Calculated MW: 748 aa, 86 kDa AG19545

> Observed MW: 86 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: Human, Mouse, Rat

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

The JAK/STAT pathway is an extensive signaling pathway downstream of cytokine receptors. STATs are cytosolic proteins with a common structure consisting of an N-terminal oligomerization domain, which favors formation of STAT dimers, followed by a DNA-binding domain and a C-terminal SRC homology-2 (SH2) domain, which is involved in association between STATs and receptors[PMID:22383755]. Signal Transducer and Activator of Transcription 4 (STAT4) is a transcription factor that is activated by IL-12 signaling and promotes Th1-cell differentiation and IFN- γ production [PMID:21998209].

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®555 STAT4 antibody (CL555-67568, Clone: 4A8C9) at dilution of 1:200.



1X10^6 mouse Th1 cells were intracellularly stained with 0.8 ug CoraLite®555 Anti-Human STAT4 (CL555-67568, Clone:4A8C9) (red), or 0.8 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).