For Research Use Only

## CoraLite®555-conjugated P53 Recombinant antibody

Catalog Number:CL555-80077

Featured Product




<b>Basic Information</b>	Catalog Number: CL555-80077	GenBank Accession Number: BC 003596	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 7157 UNIPROT ID: P04637 Full Name: tumor protein p53 Calculated MW: 44 kDa Observed MW: 53 kDa	CloneNo.: 2H13 Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima wavelengths: 557 nm / 570 nm
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat	Positive Controls: IF/ICC : HepG2 cells,	
Background Information	TP53, also known as P53 and NY-CO-13, belongs to the p53 family and has 9 isoforms. In SDS-Page, the observed molecular weight is about 53 kDa. TP53 acts as a tumor suppressor in many tumor types, including growth arrest or apoptosis depending on the physiological circumstances and cell types. It is involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. TP53 Localizes in the nucleus in most cells but found in the cytoplasm in some cells. (PMID: 26166714; PMID: 25225161)		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Procli Aliquoting is unnecessary for -20°C s	n300, 0.5% BSA, pH 7.3.	nt.


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®555 P53 antibody (CL555-80077, Clone: 2H13 ) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human P53 (CL555-80077, Clone:2H13) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).