For Research Use Only

CoraLite®555-conjugated Phospho-Caspase 9 (Ser196) Recombinant antibody



Purification Method:

Catalog Number: CL555-80346

Basic Information

Catalog Number: GenBank Accession Number:

CL555-80346 BC002452 Protein A purification GeneID (NCBI): CloneNo.:

3P16 100ul, Concentration: 1000 ug/ml by 842 Nanodrop:

UNIPROT ID: Excitation/Emission maxima Source: P55211 wavelengths: 557 nm / 570 nm Rabbit Full Name:

Isotype: caspase 9, apoptosis-related cysteine

IgG peptidase

Calculated MW: 46 kDa Observed MW: 46 kDa, 35 kDa

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human, mouse

Background Information

Caspase 9 also name as MCH6, APAF3, APAF-3, ICE-LAP6 and CASPASE-9c, is a member of the cysteine-aspartic acid protease (caspase) family. It's synthesized as a 46kDa precursor protein which can be cleaved into a 35kDa subunit and a 11kDa subunit. The phosphorylated type can be detected at 55kDa and 35kDa. It plays a central role in the mitochondrial or intrinsic apoptotic pathway that is engaged in response to many apoptotic stimuli. Once activated, caspase-9 cleaves and activates the effector caspases 3 and 7 to bring about apoptosis. It can be phosphorylated by PKB/AKT1 at Ser196, this modification will downregulate its activity and decrease apoptosis. Akt phosphorylation site found in human caspase 9 is absent in mouse caspase 9.1t's reported that there is an increase in caspase 9 expression and activity in the hypoxic brain. Inhibition of Caspase 9 activity would render opportunity to treat neurological diseases such as stroke, neurodegenerative diseases or brain injury caused by hypoxia. (PMID: 19788417, PMID: 10529400, PMID: 9812896, PMID: 18840507)

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



1X10^6 HeLa cells untreated (dashed lines) or Calyculin A treated HeLa cells were intracellularly stained with 0.25 ug CoraLite®555 Anti-Human Phospho-Caspase 9 (Ser196) (CL555-80346, Clone:3P16) (red), or 0.25 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 90% MeOH.